

Auburn University Community Practice Vaccination Protocol

Canine Vaccination Guidelines

CORE VACCINES	Initial Puppy Vaccinations (Less than or equal to 16 wks)	Initial Adult Vaccinations (Greater than 16 weeks)	Revaccination (Booster) Recommendations
Canine Distemper Virus (CDV) Canine Adenovirus-2 (CAV-2) Canine Parvovirus (CPV) Given together as DA2P, Modified Live vaccine (MLV)	All puppies should receive a minimum of 3 doses between the ages of 6 and 16 weeks, administered at intervals of 3-4 Weeks (e.g. 8, 12, and 16 weeks) The final dose should be administered at 15-16 weeks of age.	Two doses, 3-4 weeks apart	All puppies should receive a 1-year booster vaccination 1 year after completion of the initial puppy series. Following this vaccination, revaccination is recommended at intervals of every 3 years.
Rabies (RV) Killed	All puppies should receive one dose at 12 weeks* of age	Administer a single dose	Second rabies vaccination is given 1 year following administration of the initial dose, regardless of the animal's age at the time the first dose was administered. Booster vaccines should be administered every 3 years.

NONCORE VACCINES

Leptospira interrogans (L4) (With other serovars) Killed bacterin Can be given in combo with Distemper, Adeno-2, Parvo and Parainfluenza as DA2PPL (MLV w/killed Bacterin)	Administer one dose at 12 weeks and one dose 3-4 weeks later. We recommend this to all puppies in our area due to the risk of exposure. For optimal response, do not administer to puppies younger than 12 weeks.	Two doses, 2-4 weeks apart.	Annually It should be considered that dogs at exceptionally high risk should be vaccinated at 12 and 16 weeks of age, then at intervals of 6-9 months Until risk has been reduced.
Bordetella bronchiseptica and Parainfluenza (MLV) Intranasal (used primarily for faster protection)	Administer a single dose as early as 6 weeks of age. Recommended when determined to be necessary based on lifestyle of the dog.	Single dose recommended	Annually. More often in very high-risk animals. Only recommended when necessary based on lifestyle. Should be given a minimum of 72 hours before expected exposure.
Canine Influenza Virus (CIV) Killed	Administer one dose at 12 weeks and one dose at 16 weeks. Only recommended when determined to be necessary based on lifestyle of the dog.	Two doses, 2-4 weeks apart	Annually. Only recommended when determined to be necessary based on lifestyle of the dog. Vaccine does not prevent disease but decreases clinical signs.

* In accordance with Alabama Code 3:7A1-15 Section 2.(a)

Feline Vaccination Guidelines

CORE VACCINES	Initial Kitten Vaccinations (Less than or equal to 16 wks)	Initial Adult Vaccination (Greater than 16 weeks)	Revaccination (booster) Recommendations
Feline Viral Rhinotracheitis (Feline Herpesvirus-1) and Feline Calicivirus (FCV) Panleukopenia Virus (FPV) Given as combination FVRCP Modified live vaccine (MLV)	All kittens should receive 3 doses between the ages of 6 and 16 weeks of age administered at intervals of 3-4 weeks (i.e. 8, 12 and 16 weeks). The final dose should be administered at 15-16 wks of age.	Two doses, 3-4 weeks apart	A single dose is given 1 year following the last dose of the initial series. Following this vaccination, revaccination is recommended at 3 year intervals.
Rabies (RV) Recombinant (rRabies) Non-adjuvanted	All kittens should receive a single dose at 12 weeks* of age	Administer a single dose	Annual booster is required when using recombinant product
NONCORE VACCINES			
Feline Leukemia Virus (rFeLV), recombinant Non-adjuvanted Transdermal-canarypox vectored	Administer an initial dose as early as 8 weeks, booster 3-4 weeks later All kittens should be FeLV tested and confirmed negative before initial vaccination. Recommended for all kittens, Regardless of intended lifestyle.	2 doses, 3-4 weeks apart FeLV tested and negative before vaccination	Annual booster is recommended when determined to be necessary based on lifestyle of the cat. Cats should be tested yearly prior to vaccination.

* In accordance with Alabama Code 3:7A1-15 Section 2.(a)